
WAX KA BARO COVID-19 LEARN ABOUT COVID-19

Waa maxay calaamadaha lagu garto?





Qufac Cough

Murug xanuun Muscle Aches

Neefsashada oo dhib ah Trouble Breathing

Madax xanuun Headache

Daal dareemaya

Feeling Tired

Sore Throat

Dhadhanka iyo urka oo taga Loss of smell or taste

Gariir ama ilmo Shaking or Chills

Maxaa la samayn karaa si looga hortago COVID-19? What can be done to help prevent COVID-19?

Xiro maaska,ka fogoow 6 feet dadka kale markaad banaanka u baxdo, gacmaha dhaq,ka ilaali taabashada wejiga, guriga joog haddii aad jirran tahay, iska ilaali meelaha dadku ku badan yahay sida:

magaaxiyaha

🛋 Masjidka, iwm.

Wear your mask, social distance - keep 6 feet from others, wash hands, avoid touching face, stay home if sick, avoid crowded spaces such as

帶 restaurants

gathering places

🛋 mosque, etc.

Hadad isku aragto calaamadaha COVID-19: Do you have COVID-19 Symptoms?

Iska baadh COVID-19. Get a COVID-19 test.

Ka fogow dadka kale, ugu yaraan 5 maximood ilaa intaad aad ka boqsanayso. Isolate from others.

Guriga joog ilaa aad ka fiicnsato. Riest at Ixome until you are better.

> Cab cabitaanno badan oo dareereyaal ah Drink plenty of fluids.



Colorado Clinical and Translational Sciences Institute (CCTSI)
UNIVERSITY OF COLORADO DENVER | ANSCHUTZ MEDICAL CAMPUS

Ma xanuunsan tahay in ka badan 5-7 maaimood?

Are you sick longer than 5-7 days?

Haddii aad weli bukto 5-7 maalmood kadib markii laga helay COVID-19 ama ay kugu adag tahay neefsashada, la xiriir dhakhtarkaaga ama qolka degdega ee isbitaalka.

If you are still sick 5-7 days after positive COVID-19 test or you have trouble breathing, get help from primary care or hospital emergency room.

Haddii aad hesho daryeel caafimaad ama adeegyada degdegga ah ee daawaynta COVID-19ka caymisna aadan ku jirin, waxaa kaa bixin doona kharashaadka daawaynta COVID-19 barnaamijyo federaal ah.

If you receive primary care or emergency services for COVID-19 symptoms and are uninsured, federal programs will pay for COVID-19 treatment.



CO-CEAL COVID-19 Calorado Al lanca

WAX KA BARO TALAALKA COVID-19 LEARN ABOUT THE COVID-19 VACCINE

jiifiyo isbitaalka ama dhimashada COVID-19 awgeed; wayna yar tahay inaad faafiso fayraska ama aad dadka u gudbiso.

Vaccines are safe and can save you. There is a chance you could still get COVID-19 after being vaccinated, but vaccination greatly lowers your chance of hospitalization or death due to COVID-19; and you are less likely to spread the virus to those around you.



Faahfaahin ku saabsan Talaalka iyo Xoojintiisa

Vaccine & Booster Details

Talaalka Pfizer wuxu kaa ilaalinaya ama kahortagaa iin ka bedan. 90% xanuunka daran ee COVID-19

> The Pfizer vaccine gives more than 90% protection against severe COVID-19

Talaalka Modema waxa uu kahortagaa in ka badan 90% in Bad ku xanuunsato COVID-19.

The Moderna vaccine gives more than 90% protection against getting sick from COVID-19

your body over time.

Talaalka Johnson & Johnson waxa uu kahortagaa in ka badan 70% in and ku xanuunsato COVID-19.

The Johnson & Johnson vaccine gives more than 70% protection against getting slok from COVID-19

- Booster doseka waxay xoojinaysaa talaalka sababtoo ah dhammaan talaalada A booster dose is needed because all vaccines lose their ability to protect COVID-19 ka awooddooda ay ku difaacaan jirkaaga way yaraanaysaa in muda ah ka dib.
- Waxaad ka qaadan kartaa booster dose kaaga mid ka mid ah saddexda shirkadood ee talaalka.
- Talaalka Pfizer waa loo ogolaaday da'da 5 iyo ka weyn waana 2 talalal, waxaa kaloo loo ogolaaday xoojinta talaalka (booster shot) 5 bilood ka dib. The Pfizer vaccine is authorized for ages 5 and up and is 2 shots AND a booster shot 5 months later.



3 toddobaad / 3 Weeks



5 bilood / 5 Months



You can get your booster dose from any of the three vaccines companies.

Talaalka Moderna waa la oggolaaday da'da 12 iyo ka weyn waana 2 talaal, waxaa kaloo loo ogglaaday xoojinta talaalka (booster shot) 5 bilood ka dib. The Moderna vaccine is authorized for ages 12 and up and is 2 shots AND a booster 5 months later.



4 toddobaad / 4 Weeks

AND you should get a booster 2 months later.



5 bilood / 5 Months



Talaalka Jannsen (Johnson iyo Johnson) waa la oggolaaday da'da 18 iyo ka weynba waana 1 tallaal bilowga, waana inaad qaadaataa

xoojiyaha talalka 2 bilood ka dib. The Jannsen (Johnson and Johnson) vaccine is authorized for ages 18 and up and is 1 initial shot



2 bilood 2 Months



Xaqiiqooyinka COVID-19: COVID-19 FACTS:

- Talaalada COVID-19 ka waa badbaado kuma keenayaan cudurka kansar, qalal, dhicin, dhalmo la'aan, ama dhimasho; kumana siin ono fayraska COVID-19.
- Sida laga soo xigtay Xarumaha Xakamaynta iyo Kahortagga
- Cudurada ee loo yaqaano (CDC), dhammaan talaalada COVID-19 waxay la kulmeen heerar ugu sareeya ee badbaadada, waxtarka, Iyo tayada. Mid kasta oo ka mid ah talaalada COVID-19 dhamaantood waxay badbaado to
- Talaalka COVID-19 kaama liaalin doono inaad qaado hargabka. ama qabowga. Waxaad well u baahan doontaa inaad hesho labada talaal ee COVID-19 iyo tallaka hargabka.
- The COVID-19 vaccines are safe and will not cause you to get cancer, stroke, miscarriage, infertility, or death; nor will it give you
- According to the CDC, all of the COVID-19 vaccines met high standards for safety, effectiveness, and quality. Any of the COVID-19 vaccines are all safe to use in pregnancy.
- The COVID-19 vaccine will not protect you from getting the flu or common cold. You will still need to get both the COVID-19 and the